

Get natural-looking results that *last up to 1 year.**

Talk to your aesthetic specialist today.

JUVÉDERM VOLBELLA® XC Important Information APPROVED USES

JUVÉDERM VOLBELLA® XC injectable gel is for injection into the lips for lip augmentation and for correction of perioral lines in adults over the age of 21.

IMPORTANT SAFETY INFORMATION

Are there any reasons why I should not receive JUVÉDERM VOLBELLA® XC injectable gel?

Do not use this product if you have a history of multiple severe allergies or severe allergic reactions (anaphylaxis), or if you are allergic to lidocaine or Gram-positive bacterial proteins used in these products.

What precautions should my doctor advise me about?

- Tell your doctor if you are pregnant or breastfeeding. The safety of JUVEDERM VOLBELLA® XC for use during pregnancy or while breastfeeding has not been studied
- The safety of JUVÉDERM VOLBELLA® XC in patients under 22 years has not been studied
- The safety and effectiveness of JUVÉDERM VOLBELLA® XC in areas other than the lips and perioral area have not been established in controlled clinical studies

Please see additional Important Safety Information on reverse side.

FOR Lips Perioral Lines

Natural-looking results that last up to 1 year*





Actual patient. Results may vary.

Unretouched photos of paid patient taken before treatment and 1 month after treatment. A total of 1.2 mL of JUVÉDERM VOLBELLA® XC was injected into the lips.

*With optimal treatment.

IMPORTANT SAFETY INFORMATION (continued)

What precautions should my doctor advise me about? (continued)

- Tell your doctor if you have a history of excessive scarring (eg, hypertrophic scarring and keloid formations) or pigmentation disorders, as use of JUVEDERM VOLBELLA® XC may result in additional scars or changes in pigmentation
- Tell your doctor if you are planning other laser treatments or a chemical peel, as there is a possible risk of inflammation at the treatment site if these procedures are performed after treatment
- Patients who experience skin injury near the site of injection with this product may be at a higher risk for side effects
- Tell your doctor if you are on immunosuppressive therapy used to decrease the body's immune response, as use of JUVEDERM VOLBELLA® XC may result in an increased risk of infection
- Tell your doctor if you are using medications that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners, as this may result in increased bruising or bleeding at the injection site
- Minimize strenuous exercise, exposure to extensive sun or heat, and alcoholic beverages within the first 24 hours following treatment





Actual patient. Results may vary.

Unretouched photos of paid patient taken before treatment and 1 month after treatment. A total of 1.75 mL of JUVÉDERM VOLBELLA® XC was injected—0.7 mL in the perioral lines and 1.05 mL in the lips.

What are possible side effects?

The most common side effects include swelling, tenderness, bruising, firmness, lumps/bumps, redness, pain, discoloration, itching, and dryness. Most side effects are mild or moderate and last 30 days or less.

One of the risks with using this product is unintentional injection into a blood vessel, and, while rare, the complications can be serious and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring.

As with all skin injection procedures, there is a risk of infection.

To report a side effect with JUVÉDERM VOLBELLA® XC, please call Allergan at 1-800-433-8871. Please also visit Juvederm.com for more information.

Available by prescription only.



